- REST until asymptomatic
- **Graduated** return to play as directed by physician
- Follow-up with physician within 24-48 hours of injury or sooner if symptoms worsen.
- When completely recovered, written return to play release required from a health care provider trained in concussion management.



MidMichigan Medical Offices-Campus Ridge 1 4401 Campus Ridge Drive, Ste. C2000 Midland, Michigan 48640 Phone (989) 837-9350 www.midmichigan.org/wellsport

Graduated Return to Play Strategy (RTP)		
Stage	Activities	Objective
1. No activity	Complete physical and cognitive rest	Recovery
2. Light aerobic exercise	Walking, swimming or stationary cycling keeping <10% maximum predicted heart rate	Increase heart rate
3. Sport-specific exercise	Skating drills in ice hockey, running drills in soccer. No head impact activities	Add movement
4. Non-contact training drills	Progression to more complex training drills, e.g. passing drills in football and ice hockey	Exercise, coordination and cognitive load
5. Full contact practice	Following medical clearance, participate in normal training activities	Restore confidence and assess functional skills by coaching staff
6. Return to play	Normal game play	

Guidelines for Stage Progression: Each stage is 24 hours in duration. If symptoms return during a stage activity, stop all activity and rest for the entire day. The following day, return to the last stage above where symptoms did not occur and progress accordingly. Each stage should be performed symptom free before progression to the next stage.

1 OR MORE OF THESE MAY = CONCUSSION:

May not appear for **MINUTES**, **HOURS**, **DAYS** or **UNTIL CHALLENGED** (physically **OR** mentally)

Headache

Vision Changes

Mood Changes

- · Pressure in the Head
- · Sensitive to Light or Noise

Behavior or Personality Changes

- Nausea or Vomiting
- Feeling Sluggish or Groggy
- Being Knocked out (even briefly)

Sleep Changes

Confusion

Answers Questions Slowly

Dizziness

- Difficulty Concentrating or Remembering
- Moves Clumsily

Keep CONCUSSIONS on the Sidelines!

DANGER SIGNS = IMMEDIATE MEDICAL ATTENTION:

- Symptoms Get Worse
- Seizure

Neck Pain

- Decreasing Consciousness
- Vomiting

· Weakness in Arms or Legs

- Increasing Sleepiness
- Trouble Recognizing People or Places
- Slurred Speech

With proper recognition & management YOU can prevent permanent brain injury & death